

Use of waggle dance information in honey bees is linked to gene expression in the antennae, but not in the brain

Anissa Kennedy¹  | Tianfei Peng^{1,2}  | Simone M. Glaser¹ | Melissa Linn¹ |
 Susanne Foitzik¹  | Christoph Grüter^{1,3} 

¹Institute of Organismic and Molecular Evolution, Johannes Gutenberg University Mainz, Mainz, Germany

²College of Plant Science, Jilin University, Changchun, China

³School of Biological Sciences, University of Bristol, Bristol, UK

Correspondence

Anissa Kennedy, Institute of Organismic and Molecular Evolution, Johannes Gutenberg University Mainz, Biozentrum I, Hanns Dieter Hüsch Weg 15, D-55128 Mainz, Germany.

Email: anissakennedy04@gmail.com

Funding information

German Research Foundation (DFG), Grant/Award Number: DFG GR 4986/3-1 and FO 298/27-1; China Scholarship Council, Grant/Award Number: 201606170134; "Inneruniversitäre Forschungsförderung" of the University of Mainz

Abstract

Use of waggle dance information in honey bees is linked to gene expression in the antennae, but not in the brain

Decision-making in social insects relies on environmental and intrinsic factors. Honey bees use a unique form of communication, the waggle dance, to inform nestmates about the location of food sources. However, as in many other animals, experienced individuals often ignore this social information and prefer to rely on prior experiences, i.e., private information. The neurosensory factors that drive the decision to use social information are not yet understood. Here we test whether the decision to use social dance information or private information is linked to gene expression differences in different parts of the nervous system. We trained bees to collect food from sugar water feeders and observed whether they utilize social or private information when exposed to dances for a new food source. We performed transcriptome analysis of four brain parts (11–16 bees per tissue type) critical for cognition: the subesophageal ganglion, the central brain, the mushroom bodies, and the antennal lobes but, unexpectedly, detected no differences between social or private information users. In contrast, we found 413 differentially expressed genes in the antennae, suggesting that variation in sensory perception mediates the decision to use social information. Social information users were characterized by the upregulation of biogenic amine genes, while private information users upregulated several genes coding for odour perception. These results highlight that decision-making in honey bees might also depend on peripheral processes of perception rather than higher-order brain centres of information integration.

KEY WORDS

animal communication, brain, foraging, information strategies, transcriptomics, waggle dance

1 | INTRODUCTION

Exchanging information is essential in all animal societies. Communicating about resources, reproductive state, group

membership, and threats are vital in ensuring the survival and success of the group. However, relying on social information is often not the only available option, for example to find a food source, but searching for a resource individually can often be the better

Kennedy and Peng contributed equally to this work.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. Molecular Ecology published by John Wiley & Sons Ltd

choice (Dechaume-Moncharmont et al., 2005; Hoppitt & Laland, 2013; l'Anson Price et al., 2019; Kendal et al., 2009; Laland, 2004). Furthermore, an individual can rely on private information (e.g., spatial memory) about previously visited food source locations (Grüter & Leadbeater, 2014; Rendell et al., 2010). It is crucial for an organism to assess the different available options and their consequences to make the best decision in a given environment. Acquiring information through individual exploration, for instance, provides up-to-date information, but comes with the cost of trial-and-error learning. Social information avoids the costs of individual learning and exploration, but can involve the inefficient or erroneous transmission of information (Dechaume-Moncharmont et al., 2005; Giraldeau et al., 2002; l'Anson Price et al., 2019; Rieucau & Giraldeau, 2011). Thus, animals often employ flexible strategies for deciding between social or private information (Grüter & Leadbeater, 2014; Hoppitt & Laland, 2013; Kendal et al., 2009; Laland, 2004).

Social insects employ various methods to send signals to nestmates. Information exchange regarding resources is particularly well studied and a wide range of communication behaviours are used, such as tandem running in ants (Alleman et al., 2019; Glaser & Grüter, 2018; Möglich et al., 1974) and trail pheromones in ants, and stingless bees (Czaczkes et al., 2015; Hölldobler & Wilson, 2009; Jarau, 2009). Honey bees (*Apis*) use a unique form of communication, the waggle dance that gives spatial information to nestmates about both distance and direction of a food source or a nest site in relation to the sun (von Frisch, 1967). In foraging, dances are performed by returning foragers as an advertisement for high quality food sources. Furthermore, waggle dancers emit floral odours and a blend of hydrocarbons that provide additional information and stimulate foraging in unemployed foragers (Farina et al., 2012; Gilley et al., 2012; Thom et al., 2007). Only a relatively small percentage of waggle dance followers use dance information to discover new food sources. The majority of waggle dances trigger experienced foragers to resume foraging at already familiar food sources, disregarding social dance information for private spatial information (Biesmeijer & Seeley, 2005; Grüter et al., 2008). While various factors, like experience (Biesmeijer & Seeley, 2005; Grüter & Ratnieks, 2011; Richter & Waddington, 1993) and age (Tofilski, 2009; Woyciechowski & Moroń, 2009) are likely to affect whether a bee uses social information, still little is known about the neuronal basis of dance communication and its use (Barron & Plath, 2017).

Social insect behaviour and, in particular, foraging strategies are linked to gene expression in the brain (Ingram et al., 2011; Robinson et al., 2008; Toth & Robinson, 2009; Toth et al., 2010; Zayed & Robinson, 2012). Behavioural variation among workers and within foragers seems to be strongly connected to biogenic amine signalling, such as dopamine, octopamine, tyramine, and serotonin signalling (Arenas et al., 2020; Barron et al., 2002; Liang et al., 2012; Linn et al., 2020; Mercer & Menzel, 1982; Peng et al., 2020, 2021; Scheiner et al., 2002, 2017; Schulz et al., 2003). Transcriptomic differences between behavioural groups were typically studied by

investigating the entire brain (e.g., Alleman et al., 2019; Liang et al., 2012; Whitfield et al., 2003). However, different brain parts serve specific functions and are expected to differ in gene expression. For example, the antennal lobes receive input from the olfactory sensory neurons in the antennae (Paoli et al., 2016) and process olfactory information (Homberg et al., 1989; MaBouDi et al., 2017; Paoli et al., 2016). Insect mushroom bodies are the centre of multimodal sensory integration, learning and memory (Collett & Collett, 2018; Strausfeld et al., 2009), whereas the central brain supports foraging behaviour via motor control (Hanesch et al., 1989). Barron and Plath (2017) have suggested that the central brain might play a crucial role in the decoding of waggle dance information. Finally, the suboesophageal ganglion mediates reward and taste perception (Dacks et al., 2005; Kreissl et al., 1994; Sinakevitch et al., 2005). However, whether and how these different parts of the nervous system are involved in dance communication and information-use is not well understood. Furthermore, we still know little about the role of the peripheral nervous system for decision-making and information processing (see e.g., Ozaki et al., 2005). The antennae, in particular, play important functions in social insect behaviour, both within and outside the colony, such as mediating pheromone signalling (Grozinger et al., 2003; Nagari & Bloch, 2012; Pankiw et al., 2004; Vergoz et al., 2009), nestmate recognition (Ozaki et al., 2005; van Zweden & D'Ettorre, 2010) and odour learning (Robertson & Wanner, 2006; Rogers & Vallortigara, 2008).

Here we compared the gene expression of bees that used dance information (social information, SI) with those that preferred private information (PI) in different brain areas and the antennae in the honey bee *Apis mellifera*. We trained cohorts of workers to sucrose solution feeders and, subsequently, confronted them with conflicting social information about a new high-quality food source. As was shown for scouts, i.e., foragers that search for new food sources independently (Liang et al., 2012), we predicted that there are distinct neurogenomic signatures underlying the decision to use either social or private information. We compared different brain and peripheral chemosensory areas in both types of bees. We demonstrate that bees that decode and use waggle dance information differ in gene expression only in the antennae and provide evidence for roles of biogenic amine signalling and olfactory perception.

2 | MATERIALS AND METHODS

2.1 | Colony set-up

A total of six observation hives of *Apis mellifera carnica* were studied from August through October 2016 (H1–H3) and 2018 (C1–C3), each containing approximately 2000–3000 workers of mixed ages. Colonies were established from the Johannes Gutenberg University apiary in Mainz, Germany, a few weeks prior to the start of experiments. Each of the observation colonies contained three frames, brood, food reserves and were headed by a naturally mated queen.

2.2 | Training

Training was conducted one colony at a time. Workers were trained according to standard training procedures to collect sucrose solution at one of two artificial feeders (Linn et al., 2020; von Frisch, 1967). First, a cohort of 50–60 workers was trained to the training feeder (T.F.). These workers were used as the samples that would later be designated as either social or private information users on test day. Then, a smaller cohort of ~20 foragers was trained to the dance feeder (D.F.). These workers would be designated as dancers on test day. Both feeders were 150 m from the observation colonies with ca. 300 m separating the training and dance feeder (Figure S1). Workers were trained to their respective feeder with an unscented 0.8 M sucrose solution and were individually marked with a number tag on the thorax. This spatial arrangement ensured that workers would visit only one feeder and no mixing of individuals between dance and training feeders occurred. The day after training, the sucrose solution was reduced to 0.3 M at both feeders with the addition of an identical scent (5 µl of essential oil/100 ml sucrose solution). This concentration made sure that trained foragers would return to their respective feeder, but not recruit additional bees. Colonies were trained to a different odour: C1, H1 = sage, C2, H2 = jasmine, C3, H3 = peppermint. Since the same feeder locations were used throughout the experiments, a colony-specific odour was assigned to reduce the risk that for the duration of training and testing of a colony, bees from a previously studied colony would discover the feeders. During 60 min, workers were allowed to visit their feeder repeatedly (2016: 5.24 ± 3.79 visits, $N = 191$; 2018: 8.09 ± 5.17 visits, $N = 102$). The 60 min training with scented solution allowed workers to associate reward, scent, and location of the respective feeder.

2.3 | Sample collection

On the test day, the day after the 60 min odour training, 2 M sucrose solution with the same scent as used during training was offered only at the dance feeder location, while the training feeder was empty. The sucrose concentration at the dance feeder was high to induce the collecting foragers to perform waggle dances. T.F. trained workers could then decide whether to use social information by following the waggle dances performed by the returning dancers (fly to the D.F.) or disregard the dance vector information and use private information (return to the T.F.). The arrival time and capture time of each individual bee was recorded. Dance and dance following behaviour were recorded in the observation colony using a high definition camera to quantify dance following behaviour by T.F. foragers. Workers trained to the T.F. that arrived at the D.F. location were collected in Eppendorf tubes and immediately preserved in liquid nitrogen; these workers were the social information users. Workers trained to the T.F. feeder that arrived at the T.F. feeder location were collected at a similar time; these workers were the private information users.

2.4 | Video analysis

Videos were analysed using VLC Media Player. Dances and dance following behaviours were analysed frame by frame. A worker was only counted as following a dance when she was within one antennal length of a marked dancer during the waggle run phase (Grüter et al., 2008; Linn et al., 2020), which is the component of the waggle dance that encodes the vector information (von Frisch, 1967). We compared the dance following behaviour of private and social information users with linear mixed-effects models (LME and GLMMs). The nlme-package and linear mixed-effects models (LMEs) were used when the response variable was normally distributed (waggles per dance followed, averaged for each bee). The lme4-package and generalized linear mixed-effects models (GLMMs) were used when the response variable had a Poisson distribution (total number of dances followed; Zuur et al., 2009). Colony-ID and year (2016 and 2018) were included as hierarchically nested random effects to account for their effects (Zuur et al., 2009).

2.5 | Brain dissection and RNA extraction

In 2016, we dissected the calyxes of the mushroom bodies and antennal lobes from 14 workers (seven social information users and seven private information users, 2–3 per colony and type). We confirmed that all social information users followed dances extensively. In 2018 we dissected central brains and subesophageal ganglia from 16 workers (eight social information users and eight private information users, 2–3 per colony and type), and the antennae from 11 different workers (1–4 per colony and type; see Figure 1 in Sen Sarma et al., 2009 for a schematic representation of the brain areas and cutoff lines). The additional handling of the samples after being flash frozen in liquid nitrogen caused the antennae of some samples to be brittle and easily break apart. Different workers were used to ensure that whole antennae could be used for equal extraction of RNA from all samples.

Heads from individual workers were cut from the body and fixed on melted dental wax in a prechilled petri dish over ice. The antennae were cut off and stored in 100 µl of TRIzol (Invitrogen). Incisions were made at the antennal base, around the eyes, through the compound eye, and the ocellus. The cuticles, glands, retina and tissue around the brain were removed and the exposed tissues of the head were submerged with cooled bee saline (154 mM NaCl, 2 mM NaH₂PO₄, 5.5 mM Na₂HPO₄, pH 7.2). Subesophageal ganglion and central brain (which included the mushroom body peduncles, the bundled axons from the Kenyon cells in the calyxes), were removed by cutting off optic lobes, antennal lobes, and mushroom body calyxes. All tissues called “mushroom body” refer to mushroom body calyxes as it is extremely difficult to remove mushroom body peduncles. The calyxes contain the intrinsic Kenyon cells, where a large part of mushroom body transcription takes place and, therefore, the calyxes are often used to study




FIGURE 1 Principal component analysis (PCA) plots displaying variance between individual samples based on all genes for each tissue type. Samples are organized by colour according to information use strategy (blue = social, red = private) and shapes by colony ID (circles = colony 1, triangles = colony 2, squares = colony 3)

mushroom body gene expression (Humphries et al., 2003; Reim & Scheiner, 2014; Sarma et al., 2009). Furthermore, the tissue called “central brain” refers to a brain region that also includes the mushroom body peduncles and putative differences in expression in this tissue should be interpreted carefully because of the different functions of these tissues. Each dissection was completed in less than 5 min to prevent RNA degradation. Brain parts were stored in 100 ml of Trizol (Invitrogen) in -80°C for later RNA extraction using RNeasy Mini Extraction Kit (Qiagen) according to the manufacturer's protocol.

2.6 | Transcriptome analysis

For sequencing, aliquots of RNA from private and social information users were sent to Beijing Genomics Institute (BGI) for library construction and sequencing. In 2016, Hiseq 4000 was used to sequence 100 base pair (bp) paired-end reads, obtaining 40 Mio clean reads per sample. The total sample size was 28. In 2018, BGISEQ was used to sequence 100 bp paired-end reads, obtaining 70 Mio clean reads per sample. The sequencing failed for one sample and one sample was damaged during the travel (Eppendorf tube burst), decreasing our total sample size to 41. Raw reads were quality checked using FASTQC v.0.11.8 (Andrews, 2010) followed by Illumina adapter removal using TRIMOMATIC v.0.38. (Bolger et al., 2014). Clean reads were aligned using HISAT2 v.2.1.0 (Kim et al., 2017) to the honey bee genome HvA3.1 as a reference (Wallberg et al., 2019). To count

how many aligned reads mapped to genes, we used HTSEQ v.0.11.2 (Anders et al., 2015) to generate count tables. Count tables for each part were analysed separately for gene expression differences between social and private information users using the R package DESEQ2 v.1.24.0 (Love et al., 2014). Before the analysis, an additional filtering step was added to ensure that only genes with counts of at least 10 reads in at least six samples ($n - 1$ of the smallest sample size) were used in the gene expression analysis. Information strategies were compared using the likelihood ratio test (LRT) approach whereby a full model with information type (SI or PI) and colony-ID as fixed factors is compared with a reduced model containing only colony-ID, taking into consideration colony effects. Genes were considered differentially expressed if the false discovery rate (FDR) corrected p -value was $<.05$. To ensure that the number of DEGs calculated by DESeq2 were not due to chance and to account for the uneven number of samples across bee types and colonies for the antennae, we additionally performed permutations by switching samples from opposite information user groups (see methods in Libbrecht et al., 2016). For example, a sample was switched for a different information user group and the number and distribution of DEGs was compared to those calculated from our model in DESEQ2. We performed 28 permutations (14 times switching two samples for each group and 14 times switching three samples for each group) and recorded the number of DEGs in each permutation. We then compared this number to the numbers for all possible combinations of our samples to assess the number of DEGs that could be expected by chance.

We used the R package DEGREPORT v.1.20.0 (Pantano, 2019) to visualize any patterns for all genes going into the analyses and to identify clustering patterns across social and private information users by using the rlog function of DESEQ2 to generate normalized count data and the default settings. PCAs (principal components analysis) based on all genes were performed for all tissues to visualize variation between samples. A heatmap was generated to visualize the clustering patterns of the differentially expressed genes in the antenna (Figure S3). All analysis were performed in R v.3.5.0 (R Core Team, 2018).

2.7 | Gene ontology enrichment

DEGs were loaded in a BLAST search on the NCBI database against the honey bee genome HvA3.1 to find gene annotations. To further obtain information about gene ontology (GO; Ashburner et al., 2000) and KEGG pathway (Ogata et al., 1999) enrichment we used INTERPROSCAN v.5.36-75.0 (Jones et al., 2014) on the protein sequences. The R package TOPGO v.2.36.0 (Alexa and Rahnenfuhrer, 2018) was used to perform an enrichment analysis of GO terms and a Fisher's exact test was performed on the list of biological processes.

3 | RESULTS

3.1 | Dance following of social and private information users

Dance following behaviour was analysed by combining data collected from video analysis for both years. Using a linear mixed-effects model, we found that SI bees followed fewer dances than PI bees during the testing period (5 ± 0.7559 vs. 7.091 ± 1.546 dances) (GLMM: z-value = -2.122; $p = .0338$). However, SI bees followed dances for longer (more waggle runs per dance) than PI bees (27.214 ± 4.089 vs. 30.818 ± 6.5 ; LME: t-value = 2.218; $p = .0396$).

3.2 | Gene expression analysis

The likelihood ratio test (LRT) comparison of information use strategies revealed no differences in gene expression between the two information user groups in the central brain, antennal lobes, and subesophageal ganglion (Figure 1). There was only one differentially expressed gene between social and private information users from our mushroom body calyxes' samples, which encodes for an uncharacterized protein ($p = .026$, gene ID: rna-XR_003305479.1). However, there were 413 differentially expressed genes in the antennae, 318 were higher expressed in social information users and 95 were higher expressed in private information users. To confirm these substantial differences in gene expression in the antennae, we used permutations of samples to assess how this affects the number of DEGs in the antennae. The permutations showed that only very few DEGs were detected when 2–3 samples were swapped between

the SI and PI groups and their respective colonies (colony ID as fixed factor: 11.89 ± 31.87 , $N = 28$; colony ID not included: 3.25 ± 7.01 , $N = 28$; Figure S2). This confirms that the substantial differences in gene expression in the antennae are linked to whether bees belonged to the SI or the PI group. PCA plots used transformed data of all genes to further explore whether there is a clustering of samples based on information use strategies and colony. While a clustering pattern based on information use and colony can be seen for the antennae (Figure 1), the other tissues showed no clear clustering according to information use.

Exploring the list of DEGs in the antennae revealed that numerous odourant binding and chemosensory proteins differed in their expression in social and private information users. Specifically, we detected five genes for odourant or chemical perception among the upregulated genes in private information users (*odourant binding protein 5, 11, 19, 7* and *chemosensory protein 1*) and two among the upregulated genes in social information users (*odourant binding protein 7* and *chemosensory protein 2*; Figure 2). Several genes involved in biogenic amine production or signalling were also differentially expressed. Social information users had a higher expression of *tyrosine kinase Btk29A*; *dopamine N-acetyltransferase, tryptophan 5-hydroxylase 1*, which are involved in the production of dopamine or serotonin (Coleman & Neckameyer, 2005; Sasaki et al., 2012; Vavricka et al., 2014), while private information users had a higher expression of one gene *tyramine receptor, transcript variant X1*, which is associated with biogenic amine signalling (Blenau et al., 2000; Mustard et al., 2005; Figure 3). Social information users also had higher expression of the egg yolk precursor protein *vitellogenin*, a gene that is upregulated in nurses and downregulated in foragers' fat bodies and brain (Amdam et al., 2003; Nunes et al., 2013; Figure 3).

3.3 | Gene function and enrichment analysis

Separate GO enrichment analyses of only upregulated genes for each information strategy showed a small number of enriched functions: nine biological processes enriched in social information users connected mainly on carbohydrate (10 genes) and lipid (7) metabolic process and 18 enriched biological processes in private information users focused on oxidation-reduction process (7) and protein catabolic process (11).

4 | DISCUSSION

Information and its use in animals is an important topic in behaviour, ecology, and evolution because information is a critical currency that allows animals to make adaptive decisions in a given situation (e.g., Dall et al., 2005; Danchin et al., 2004; Hoppitt & Laland, 2013; Rieucau & Giralda, 2011). The decision of when to utilize social versus private information to best exploit potential opportunities while avoiding costs is crucial for success and has been studied within a variety of both social and nonsocial animals (e.g., Bonnie & Earley, 2007;




FIGURE 2 Plots of individual odourant binding protein (OBP) and chemosensory protein (CSP) genes. Black dots show counts for individual samples and shapes correspond to the colony ID (circle = colony 1, triangle = colony 2, square = colony 3). Coloured circles are representative of the average for the respective information strategy (red = private, blue = social) with confidence intervals. (a) OBP11 ($p < .001$), (b) OBP19 ($p = .001$), (c) OBP5 ($p = .03$), (d) OBP7 ($p < .001$), (e) CSP1 ($p = .009$) and (f) CSP2 ($p = .007$). p -values shown are after FDR correction

Grüter & Ratnieks, 2011; Haak et al., 2020; Taborsky & Oliveira, 2012; Weimerskirch et al., 2010; Wray et al., 2012). However, it is still unclear whether and how molecular and neurosensory factors determine an individual's preference for social or private information.

Here we explore whether gene expression differences between honey bee foragers are linked to the use of social or private information about food sources to uncover the potential molecular mechanisms that underlie the decision to decode and use waggle dances in honey bees. Contrary to our prediction, the transcriptomes of all four analysed brain parts did not differ between bees using these two foraging strategies. Strikingly, however, we found substantial gene expression differences in the antennae. Over 400 genes were differentially expressed between social and private information users, suggesting that the sensory perception of these two forager types differs. This is further supported by our results which show gene expression differences related to odourant binding proteins, chemosensory proteins, and genes associated with biogenic amine production.

The lack of differences in the brain areas was unexpected given that Liang et al. (2012) found extensive differences in whole-brain gene expression between scouts and nonscout foragers (in their study, nonscout foragers could have included both private and

social information users). We expected the mushroom bodies to show differences since it has previously been shown that they are involved in multisensory integration, learning, and place memory (e.g., Collett & Collett, 2018; Strausfeld et al., 2009). The antennal lobes are involved in odour recognition and memory through the interconnectivity of neurons with the mushroom body and were thus selected as another area of interest (Boeckh & Tolbert, 1993). The central brain has been suggested as an important area for dance communication (Barron & Plath, 2017), while the subesophageal ganglion plays important roles in reward perception and taste (Galizia et al., 2011). Together, these brain regions were thought to process reward and odour perception which could play an important role in the decision to use dance information. Our study indicates that information use strategies may not primarily depend on integration of information in higher order centres, but that the antennae play a major role in decision-making when facing communication signals. However, we note that the sample size in this study could have limited our ability to detect small differences in brain tissues.

The 413 differentially expressed genes in the antennae present an array of gene families and functions. Of particular interest




FIGURE 3 Plots of individual genes associated with biogenic amine production and reproduction. Black dots show counts for individual samples and shapes correspond to the colony ID (circle = colony 1, triangle = colony 2, square = colony 3). Coloured circles are representative of the average for the respective information strategy (red = private, blue = social) with confidence intervals. (a) Tyramine receptor ($p = .018$), (b) Vitellogenin ($p = .045$), (c) Tyrosine kinase Btk29A ($p = .006$) and (d) Tryptophan 5-hydroxylase 1 ($p < .001$). p -values shown are after FDR correction

are genes coding for odourant binding proteins and those involved in biogenic amine production and signalling due to their potential roles in chemosensory perception. Thus, differences in the perception of chemosensory information cues and signals could result in divergent foraging strategies. While our study cannot disentangle whether gene expression is the cause or the consequence of the information use strategy, they suggest that chemosensory perception by the antennae could be involved in the decision to decode waggle dances and use social information. In many social insects, the antennae play an integral role in social recognition (Balbuena & Farina, 2020; Ozaki et al., 2005; Sharma et al., 2015). Studies in *Oecophylla smaragdina*, for instance, indicated that the density of antennal sensilla is important in regulating behaviour, particularly in determining the aggression response behaviour to non-nestmates (Cholé et al., 2019; Gill et al., 2013). Similar to other social insects, honey bee foragers first use their antennae to perceive and respond to a variety of chemical signals for navigation (Menzel & Greggers, 2013), efficient nectar/pollen collection (Arenas & Farina, 2012), and dance communication (Gilley et al., 2012; Reinhard & Srinivasan, 2009; Thom et al., 2007).

By transporting odourants, for example, from antennal sensilla to odourant receptors, odourant binding proteins (OBPs) play important roles in olfactory sensitivity (Leal, 2013). They are hypothesized to be important in insect communication (Pelosi et al., 2005), including in the honey bee which use highly complex odours and pheromones to regulate their social activities (Baracchi et al., 2020; Farina et al., 2012). Of the 21 OBPs found in the honey bee, only

nine are exclusively expressed in the antennae. The remaining OBPs are active throughout the honey bee body or specific nonolfactory tissues (Forêt & Maleszka, 2006). Our analysis revealed that workers which rely on private information in the form of spatial memory show higher expression of four odourant binding proteins (*obp5*, *obp11*, *obp19*, and *obp7*), whereas workers that rely on socially acquired information upregulate one (*obp7*). Thus, ~25% of all OBPs found in honey bees were differentially expressed. Of the OBPs that were upregulated in private information users, *obp5* and *obp11* have been previously shown to be exclusively expressed in the antennae and suggest a chemosensory function (Forêt & Maleszka, 2006). Interestingly, *obp11* is mainly expressed in a rare type of antennal sensilla found only in female honey bees, the *sensilla basiconica*, and is likely to facilitate the function of these sensilla (Kucharski et al., 2016). While the ligand of *obp11* remains unknown, there is evidence that the *sensilla basiconica* play important roles in the perception of cuticular hydrocarbons (CHCs) in ants (Sharma et al., 2015) and may play a similar role in honey bees (Kucharski et al., 2016). This is remarkable because CHCs emitted by dancing bees are known to trigger the use of private information in honey bees (Gilley et al., 2012; Thom et al., 2007). This raises the possibility that a higher expression of *obp11* increases the sensitivity of bees towards CHCs emitted by waggle dancers, thereby triggering private information use. The latter could also partly explain why private information users frequently follow dances, but follow them less long than social information users (Grüter & Ratnieks 2011; this study). The remaining differentially expressed OBPs (*obp19* and *obp7*) have been shown to be ubiquitously expressed, which suggests they may have additional molecular functions which we currently do not know. Overall, these results indicate a difference in perceptual sensitivity where workers which use private information perceive some chemosensory stimuli more or differently than social information users. This could have far reaching consequences for their behaviour given the role that odours play in the decision-making and information use of a forager, for example, in the identification and learning of floral resources or the perception of cuticular hydrocarbons (Gilley et al., 2012; Grüter et al., 2008; Johnson, 1967; Reinhard et al., 2004; von Frisch, 1967).

Chemosensory proteins serve a similar role as OBPs in transporting chemical stimuli through mechanisms that are not yet well understood. These proteins are heavily concentrated in antennal sensilla but are also expressed in nonolfactory tissues (Calvello et al., 2005; Forêt et al., 2007). Of the six chemosensory proteins found in honey bees (McKenzie et al., 2014), two were differentially expressed in social and private information users, chemosensory proteins 1 and 2. Both chemosensory proteins have been shown to be highly expressed in the antennae (Li et al., 2016), which further supports the view that the differences between the information strategies may be rooted in chemoreception.

Biogenic amines have been associated with regulating learning, foraging behaviour, and the transition from in-hive tasks to foraging (Lehman et al., 2006). Biogenic amine signalling is known to change with age and tissue location in honey bees (e.g., McQuillan et al.,

2012; Perry & Barron, 2013; Reim & Scheiner, 2014; Thamm et al., 2017). Specifically, dopamine, serotonin, octopamine, and tyramine titres in the brain were found to be linked to both task and age (Barron et al., 2002; Harris and Woodring, 1992; Kokay & Mercer, 1997; Schulz & Robinson, 1999). For example, tyramine levels have been linked to novelty seeking in scouting behaviour (Cook et al., 2018; Liang et al., 2012), sucrose responsiveness (Scheiner, Entler, et al., 2017; Scheiner et al., 2002; Scheiner, Reim, et al., 2017), and division of labour between nectar and pollen foragers (Hunt et al., 1995; Scheiner et al., 2001). Dopamine has been shown to modulate sucrose responsiveness (Scheiner et al., 2002), learning (Vergoz et al., 2007) and dance following (Linn et al., 2020), whereas serotonin influences foraging activity (Schulz et al., 2003) and regulates feeding in many animals (Blundell & Halford, 1998; French et al., 2014; Voigt & Fink, 2015). Our findings of an upregulation of genes associated with biogenic amine production, raise the possibility that social information users could differ in their sensory perception as well as sucrose response thresholds compared to private information users. It is noteworthy, however, that the differences we found in relation to biogenic amine signalling were not in the brain. Instead, higher expression of several genes associated with dopamine and serotonin production was found in the antennae of social information users. We did not control for foraging age or experience, which have already been shown to affect gene biogenic amine expression (Reim & Scheiner, 2014). However, the lack of differential expression in brain areas suggests that there was no systematic age bias in our samples.

Another interesting differentially expressed gene, *vitellogenin*, is best known as an egg yolk precursor protein for egg laying organisms. Under normal conditions in social insects, the queen is the main reproductive member and therefore produces the highest levels of *vitellogenin*. However, *vitellogenin* serves important roles for other behaviours and functions outside of reproduction (Morandin et al., 2014; Nelson et al., 2007). For example, nurses produce the next highest levels of *vitellogenin* in their hypopharyngeal glands to fortify brood food with protein (Amdam et al., 2003, 2009; Wegener et al., 2009). A characteristic feature of the transition from nurse to forager is the drop in *vitellogenin* levels (Amdam et al., 2003; Messan et al., 2018). Our finding is consistent with evidence that biogenic amine levels are linked to *vitellogenin* and foraging behaviour (Koywiwatratkul et al., 2005; Linn et al., 2020), where social information have a similar physiological state to nurses.

Intrinsic factors such as genetic differences could also affect the decision to decode waggle dances. Honey bee queens can mate with more than 20 drones (Strassmann, 2001), and the patriline composition of our samples is not known. It is well known that different patrilines can differ in foraging behaviours, such as foraging age (Kolmes et al., 1989). Paternal effects can also impact gustatory responsiveness and learning abilities (Behrends et al., 2007; Scheiner and Arnold, 2010; Scheiner et al., 1999, 2001, 2005). It is unclear whether systematic patriline differences in the composition of PI and SI bees would lead to differential gene expression only in the

antennae, but future studies should explore whether bees using private or social information differ in their patrilines.

Overall, our results suggest an important role of the antennae in mediating decision-making and information use. In particular, we suggest a link between chemosensory perception and the reliance on communication in honey bees. For instance, private and social information users might perceive important social and environmental odours differently. Our results and those of Ozaki et al. (2005) highlight our limited understanding of the cognitive and neuronal processes that underlie animal decision-making. The relative importance of peripheral and central information processing for decision-making is a fascinating avenue for future research. Further studies are also needed to disentangle the potential effects of genetic differences (i.e., different patrilines), differences in foraging experience, and other factors on gene expression. In addition, we need studies such as sucrose response thresholds, odour learning, and electroantennograms to confirm our hypothesis that the gene expression differences in the antennae of SI and PI bees is a result of differences in sensory perception.

ACKNOWLEDGMENTS

We are grateful to Semi Brami for help with data collection and Romain Libbrecht and Marah Stoldt for their support in the gene expression analyses. We thank Ricarda Scheiner and Markus Thamm for helpful advice on the brain dissections. We also would like to thank the German Research Foundation (DFG) for a grant to C. G. and S. F. (DFG GR 4986/3-1 and FO 298/27-1). T.P. was supported by a fellowship of the China Scholarship Council (File No. 201606170134). S.M.G. was supported by a grant from the "Inneruniversitäre Forschungsförderung" of the University of Mainz. Our laboratory work was supported by Marion Kever and Steffi Emmling.

AUTHOR CONTRIBUTIONS

Anissa Kennedy, Tianfei Peng, Melissa Linn, Simone M. Glaser, Susanne Foitzik and Christoph Grüter conceived the study and designed the experiments. Anissa Kennedy, Tianfei Peng, Melissa Linn and Simone M. Glaser conducted the field experiments. Tianfei Peng, Melissa Linn and Simone M. Glaser performed all honey bee brain dissections and RNA extractions. Tianfei Peng and Anissa Kennedy worked together on gene expression analysis. Data analysis was supported by Susanne Foitzik and Christoph Grüter. Anissa Kennedy wrote the first draft of the manuscript, all authors contributed to writing the final version.

DATA AVAILABILITY STATEMENT

All the Supporting Information material and additional data generated and used throughout this project may be found within the Dryad repository <https://doi.org/10.5061/dryad.sqv9s4n3g>, which contains the following: dance following information for all samples, lists of differentially expressed genes for all brain tissues, InterProScan annotation of gene lists and GO enrichments from all antenna samples, and all information for permutations including design, number of genes, and gene information. All sequencing data has been

deposited in the Sequencing Read Archive (SRA: PRJNA707149) of the NCBI.

ORCID

- Anissa Kennedy  <https://orcid.org/0000-0002-2585-4366>
 Tianfei Peng  <https://orcid.org/0000-0003-2239-9967>
 Susanne Foitzik  <https://orcid.org/0000-0001-8161-6306>
 Christoph Grütter  <https://orcid.org/0000-0001-5553-5465>

REFERENCES

- Alexa, A., & Rahnenfuhrer, J. (2018). TOPGO: Enrichment analysis for gene ontology. R package version 2.28.0.
- Alleman, A., Stoldt, M., Feldmeyer, B., & Foitzik, S. (2019). Tandem running and scouting behaviour are characterized by up-regulation of learning and memory formation genes within the ant brain. *Molecular Ecology*, 28, 2342–2359. <https://doi.org/10.1111/mec.15079>.
- Amdam, G. V., Norberg, K., Hagen, A., & Omholt, S. W. (2003). Social exploitation of vitellogenin. *Proceedings of the National Academy of Sciences*, 100, 1799–1802. <https://doi.org/10.1073/pnas.0333979100>.
- Amdam, G. V., Rueppell, O., Fondrk, M. K., Page, R. E., & Nelson, C. M. (2009). The nurse's load: Early-life exposure to brood-rearing affects behavior and lifespan in honey bees (*Apis mellifera*). *Experimental Gerontology*, 44, 467–471. <https://doi.org/10.1016/j.exger.2009.02.013>.
- Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq—A Python framework to work with high-throughput sequencing data. *Bioinformatics*, 31, 166–169. <https://doi.org/10.1093/bioinformatics/btu638>.
- Andrews, S. (2010). FastQC: A quality control tool for high throughput sequence data. Retrieved from <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Arenas, A., & Farina, W. M. (2012). Learned olfactory cues affect pollen-foraging preferences in honeybees, *Apis mellifera*. *Animal Behaviour*, 83, 1023–1033. <https://doi.org/10.1016/j.anbehav.2012.01.026>.
- Arenas, A., Lajad, R., Peng, T., Grütter, C., & Farina, W. (2020). Correlation between octopaminergic signalling and foraging task specialisation in honeybees. *Genes, Brain and Behavior*, 20, e12718. <https://doi.org/10.1111/gbb.12718>.
- Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., Davis, A. P., Dolinski, K., Dwight, S. S., Eppig, J. T., Harris, M. A., Hill, D. P., Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J. C., Richardson, J. E., Ringwald, M., Rubin, G. M., & Sherlock, G. (2000). Gene ontology: Tool for the unification of biology. *Nature Genetics*, 25, 25–29. <https://doi.org/10.1038/75556>.
- Balbuena, M. S., & Farina, W. M. (2020). Chemosensory reception in the stingless bee *tetragonisca angustula*. *Journal of Insect Physiology*, 125, 104076. <https://doi.org/10.1016/j.jinsphys.2020.104076>.
- Baracchi, D., Cabriol, A., Devaud, J. M., Haase, A., d'Ettorre, P., & Giurfa, M. (2020). Pheromone components affect motivation and induce persistent modulation of associative learning and memory in honey bees. *Communications Biology*, 3, 1–9. <https://doi.org/10.1038/s42003-020-01183-x>.
- Barron, A. B., & Plath, J. A. (2017). The evolution of honey bee dance communication: A mechanistic perspective. *Journal of Experimental Biology*, 220, 4339–4346. <https://doi.org/10.1242/jeb.142778>.
- Barron, A., Schulz, D., & Robinson, G. (2002). Octopamine modulates responsiveness to foraging-related stimuli in honey bees (*Apis mellifera*). *Journal of Comparative Physiology A*, 188, 603–610. <https://doi.org/10.1007/s00359-002-0335-5>.
- Behrends, A., Scheiner, R., Baker, N., & Amdam, G. V. (2007). Cognitive aging is linked to social role in honey bees (*Apis mellifera*). *Experimental Gerontology*, 42, 1146–1153. <https://doi.org/10.1016/j.exger.2007.09.003>.
- Biesmeijer, J. C., & Seeley, T. D. (2005). The use of waggle dance information by honey bees throughout their foraging careers. *Behavioral Ecology and Sociobiology*, 59, 133–142. <https://doi.org/10.1007/s00265-005-0019-6>.
- Blenau, W., Balfanz, S., & Baumann, A. (2000). Amtyr1: Characterization of a gene from honeybee (*Apis mellifera*) brain encoding a functional tyramine receptor. *Journal of Neurochemistry*, 74, 900–908. <https://doi.org/10.1046/j.1471-4159.2000.0740900.x>.
- Blundell, J. E., & Halford, J. C. (1998). Serotonin and appetite regulation. *CNS Drugs*, 9, 473–495. <https://doi.org/10.2165/00023210-19980906-00005>.
- Boeckh, J., & Tolbert, L. P. (1993). Synaptic organization and development of the antennal lobe in insects. *Microscopy Research and Technique*, 24, 260–280. <https://doi.org/10.1002/jemt.1070240305>.
- Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics*, 30, 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bonnie, K. E., & Earley, R. L. (2007). Expanding the scope for social information use. *Animal Behaviour*, 74, 171–181. <https://doi.org/10.1016/j.anbehav.2006.12.009>.
- Calvello, M., Brandazza, A., Navarrini, A., Dani, F. R., Turillazzi, S., Felicioli, A., & Pelosi, P. (2005). Expression of odorant-binding proteins and chemosensory proteins in some hymenoptera. *Insect Biochemistry & Molecular Biology*, 35, 297–307. <https://doi.org/10.1016/j.ibmb.2005.01.002>.
- Cholé, H., Carcaud, J., Mazeau, H., Famié, S., Arnold, G., & Sandoz, J. C. (2019). Social contact acts as appetitive reinforcement and supports associative learning in honeybees. *Current Biology*, 29, 1407–1413. <https://doi.org/10.1016/j.cub.2019.03.025>.
- Coleman, C. M., & Neckameyer, W. S. (2005). Serotonin synthesis by two distinct enzymes in *Drosophila melanogaster*. *Archives of Insect Biochemistry and Physiology: Published in Collaboration with the Entomological Society of America*, 59, 12–31. <https://doi.org/10.1002/arch.20050>.
- Collett, M., & Collett, T. S. (2018). How does the insect central complex use mushroom body output for steering? *Current Biology*, 28, R733–R734. <https://doi.org/10.1016/j.cub.2018.05.060>.
- Cook, C. N., Mosqueiro, T., Brent, C. S., Ozturk, C., Gadau, J., Pinter-Wollman, N., & Smith, B. H. (2018). Individual differences in learning and biogenic amine levels influence the behavioural division between foraging honey bee scouts and recruits. *Journal of Animal Ecology*, 88, 236–246. <https://doi.org/10.1111/1365-2656.12911>.
- Czaczkes, T. J., Grütter, C., & Ratnieks, F. L. (2015). Trail pheromones: An integrative view of their role in social insect colony organization. *Annual Review of Entomology*, 60, 581–599. <https://doi.org/10.1146/annurev-ento-010814-020627>.
- Dacks, A. M., Christensen, T. A., Agricola, H. J., Wollweber, L., & Hildebrand, J. G. (2005). Octopamine-immunoreactive neurons in the brain and subesophageal ganglion of the hawkmoth *Manduca sexta*. *Journal of Comparative Neurology*, 488, 255–268. <https://doi.org/10.1002/cne.20556>.
- Dall, S. R., Giraldeau, L. A., Olsson, O., McNamara, J. M., & Stephens, D. W. (2005). Information and its use by animals in evolutionary ecology. *Trends in Ecology & Evolution*, 20, 187–193. <https://doi.org/10.1016/j.tree.2005.01.010>.
- Danchin, E., Giraldeau, L. A., Valone, T. J., & Wagner, R. H. (2004). Public information: From nosy neighbors to cultural evolution. *Science*, 305, 487–491. <https://doi.org/10.1126/science.1098254>.
- Dechaume-Moncharmont, F. X., Dornhaus, A., Houston, A. I., McNamara, J. M., Collins, E. J., & Franks, N. R. (2005). The hidden cost of information in collective foraging. *Proceedings of the Royal Society B: Biological Sciences*, 272, 1689–1695. <https://doi.org/10.1098/rspb.2005.3137>.

- Farina, W. M., Grüter, C., & Arenas, A. (2012). Olfactory information transfer during recruitment in honey bees. In G. C. Galizia, D. Eisenhardt, & M. Giurfa (Eds.), *Honeybee neurobiology and behavior* (pp. 89–101). Springer.
- Forêt, S., & Maleszka, R. (2006). Function and evolution of a gene family encoding odorant binding-like proteins in a social insect, the honey bee (*Apis mellifera*). *Genome Research*, 16, 1404–1413. <https://doi.org/10.1101/gr.5075706>.
- Forêt, S., Wanner, K. W., & Maleszka, R. (2007). Chemosensory proteins in the honey bee: Insights from the annotated genome, comparative analyses and expressional profiling. *Insect Biochemistry & Molecular Biology*, 37, 19–28. <https://doi.org/10.1016/j.ibmb.2006.09.009>.
- French, A. S., Simcock, K. L., Rolke, D., Gartside, S. E., Blenau, W., & Wright, G. A. (2014). The role of serotonin in feeding and gut contractions in the honeybee. *Journal of Insect Physiology*, 61, 8–15. <https://doi.org/10.1016/j.jinsphys.2013.12.005>.
- Galizia, C. G., Eisenhardt, D., & Giurfa, M. (2011). *Honeybee neurobiology and behavior: A tribute to Randolph Menzel*. Springer.
- Gill, K. P., van Wilgenburg, E., Macmillan, D. L., & Elgar, M. A. (2013). Density of antennal sensilla influences efficacy of communication in a social insect. *The American Naturalist*, 182, 834–840. <https://doi.org/10.1086/673712>.
- Gilley, D. C., Kuzora, J. M., & Thom, C. (2012). Hydrocarbons emitted by waggle-dancing honey bees stimulate colony foraging activity by causing experienced foragers to exploit known food sources. *Apidologie*, 43, 85–94. <https://doi.org/10.1007/s13592-011-0080-2>.
- Giraldeau, L. A., Valone, T. J., & Templeton, J. J. (2002). Potential disadvantages of using socially acquired information. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 357, 1559–1566. <https://doi.org/10.1098/rstb.2002.1065>.
- Glaser, S. M., & Grüter, C. (2018). Ants (*Temnothorax nylanderi*) adjust tandem running when food source distance exposes them to greater risks. *Behavioral Ecology and Sociobiology*, 72, 40. <https://doi.org/10.1007/s00265-018-2453-2>.
- Grozinger, C. M., Sharabash, N. M., Whitfield, C. W., & Robinson, G. E. (2003). Pheromone-mediated gene expression in the honey bee brain. *Proceedings of the National Academy of Sciences*, 100, 14519–14525. <https://doi.org/10.1073/pnas.2335884100>.
- Grüter, C., Balbuena, M. S., & Farina, W. M. (2008). Informational conflicts created by the waggle dance. *Proceedings of the Royal Society B: Biological Sciences*, 275, 1321–1327. <https://doi.org/10.1098/rspb.2008.0186>.
- Grüter, C., & Leadbeater, E. (2014). Insights from insects about adaptive social information use. *Trends in Ecology & Evolution*, 29, 177–184. <https://doi.org/10.1016/j.tree.2014.01.004>.
- Grüter, C., & Ratnieks, F. L. (2011). Honeybee foragers increase the use of waggle dance information when private information becomes unrewarding. *Animal Behaviour*, 81, 949–954. <https://doi.org/10.1016/j.anbehav.2011.01.014>.
- Haak, C. R., Hui, F. K., Cowles, G. W., & Danylchuk, A. J. (2020). Positive interspecific associations consistent with social information use shape juvenile fish assemblages. *Ecology*, 101, e02920. <https://doi.org/10.1002/ecy.2920>.
- Hanesch, U., Fischbach, K. F., & Heisenberg, M. (1989). Neuronal architecture of the central complex in *Drosophila melanogaster*. *Cell and Tissue Research*, 257, 343–366. <https://doi.org/10.1007/BF00261838>.
- Harris, J. W., & Woodring, J. (1992). Effects of stress, age, season, and source colony on levels of octopamine, dopamine and serotonin in the honey bee (*Apis mellifera* L.) brain. *Journal of Insect Physiology*, 38, 29–35. [https://doi.org/10.1016/0022-1910\(92\)90019-A](https://doi.org/10.1016/0022-1910(92)90019-A).
- Hölldobler, B., & Wilson, E. O. (2009). *The superorganism: The beauty, elegance, and strangeness of insect societies*. WW Norton & Company.
- Homberg, U., Christensen, T. A., & Hildebrand, J. G. (1989). Structure and function of the deutocerebrum in insects. *Annual Review of Entomology*, 34, 477–501. <https://doi.org/10.1146/annurev.en.34.010189.002401>.
- Hoppitt, W., & Laland, K. N. (2013). *Social learning: An introduction to mechanisms, methods, and models*. Princeton University Press.
- Humphries, M. A., Mustard, J. A., Hunter, S. J., Mercer, A., Ward, V., & Ebert, P. R. (2003). Invertebrate D2 type dopamine receptor exhibits age-based plasticity of expression in the mushroom bodies of the honeybee brain. *Journal of Neurobiology*, 55, 315–330. <https://doi.org/10.1002/neu.10209>.
- Hunt, G. J., Page, R. E., Fondrk, M. K., & Dullum, C. J. (1995). Major quantitative trait loci affecting honey bee foraging behavior. *Genetics*, 141, 1537–1545. <https://doi.org/10.1002/gcc.2870140411>.
- I'Anson Price, R., Dulex, N., Vial, N., Vincent, C., & Grüter, C. (2019). Honeybees forage more successfully without the "dance language" in challenging environments. *Science Advances*, 5, eaat0450. <https://doi.org/10.1126/sciadv.aat0450>.
- Ingram, K. K., Kleeman, L., & Peteru, S. (2011). Differential regulation of the foraging gene associated with task behaviors in harvester ants. *BMC Ecology*, 11, 1–9. <https://doi.org/10.1186/1472-6785-11-19>.
- Jarau, S. (2009). Chemical communication during food exploitation in stingless bees. In S. Jarau, & M. Hrncir (Eds.), *Food exploitation by social insects: Ecological, behavioral, and theoretical approaches* (pp. 223–249). CRC Press.
- Johnson, D. L. (1967). Communication among honey bees with field experience. *Animal Behaviour*, 15, 487–492. [https://doi.org/10.1016/0003-3472\(67\)90048-6](https://doi.org/10.1016/0003-3472(67)90048-6).
- Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesceat, S., Quinn, A. F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., & Hunter, S. (2014). InterProScan 5: Genome-scale protein function classification. *Bioinformatics*, 30, 1236–1240. <https://doi.org/10.1093/bioinformatics/btu031>.
- Kendal, J., Giraldeau, L. A., & Laland, K. (2009). The evolution of social learning rules: Payoff-biased and frequency-dependent biased transmission. *Journal of Theoretical Biology*, 260, 210–219. <https://doi.org/10.1016/j.jtbi.2009.05.029>.
- Kim, D., Langmead, B., & Salzberg, S. (2017). HISAT2: Graph-based alignment of next-generation sequencing reads to a population of genomes. Johns Hopkins University. Center for Computational Biology. <https://doi.org/10.1101/012591>.
- Kokay, I. C., & Mercer, A. R. (1997). Age-related changes in dopamine receptor densities in the brain of the honey bee, *Apis mellifera*. *Journal of Comparative Physiology A*, 181, 415–423. <https://doi.org/10.1007/s003590050125>.
- Kolmes, S. A., Winston, M. L., & Fergusson, L. A. (1989). The division of labor among worker honey bees (Hymenoptera: Apidae): The effects of multiple patrilines. *Journal of the Kansas Entomological Society*, 62, 80–95. [https://doi.org/10.1016/0022-1910\(89\)90027-9](https://doi.org/10.1016/0022-1910(89)90027-9).
- Koywiwatratkul, P., Thompson, G. J., Sitthipraneed, S., Oldroyd, B. P., & Maleszka, R. (2005). Effects of carbon dioxide narcosis on ovary activation and gene expression in worker honeybees, *Apis mellifera*. *Journal of Insect Science*, 5, 1–10. <https://doi.org/10.1093/jis/5.1.36>.
- Kreissl, S., Eichmüller, S., Bicker, G., Rapus, J., & Eckert, M. (1994). Octopamine-like immunoreactivity in the brain and subesophageal ganglion of the honeybee. *Journal of Comparative Neurology*, 348, 583–595. <https://doi.org/10.1002/cne.903480408>.
- Kucharski, R., Maleszka, J., & Maleszka, R. (2016). A possible role of DNA methylation in functional divergence of a fast evolving duplicate gene encoding odorant binding protein 11 in the honeybee. *Proceedings Biological Sciences*, 283, 20160558. <https://doi.org/10.1098/rspb.2016.0558>.
- Laland, K. N. (2004). Social learning strategies. *Animal Learning & Behavior*, 32, 4–14. <https://doi.org/10.3758/BF03196002>.

- Leal, W. S. (2013). Odorant reception in insects: Roles of receptors, binding proteins, and degrading enzymes. *Annual Review of Entomology*, 58, 373–391. <https://doi.org/10.1146/annurev-ento-120811-153635>.
- Lehman, H. K., Schulz, D. J., Barron, A. B., Wraith, L., Hardison, C., Whitney, S., & Robinson, G. E. (2006). Division of labor in the honey bee (*Apis mellifera*): The role of tyramine β -hydroxylase. *Journal of Experimental Biology*, 209, 2774–2784. <https://doi.org/10.1242/jeb.02296>.
- Li, H. L., Ni, C. X., Tan, J., Zhang, L. Y., & Hu, F. L. (2016). Chemosensory proteins of the eastern honeybee, *Apis cerana*: Identification, tissue distribution and olfactory related functional characterization. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 194, 11–19. <https://doi.org/10.1016/j.cbpb.2015.11.014>.
- Liang, Z. S., Nguyen, T., Mattila, H. R., Rodriguez-Zas, S. L., Seeley, T. D., & Robinson, G. E. (2012). Molecular determinants of scouting behavior in honey bees. *Science*, 335, 1225–1228. <https://doi.org/10.1126/science.1213962>.
- Libbrecht, R., Oxley, P. R., Keller, L., & Kronauer, D. J. C. (2016). Robust DNA methylation in the clonal raider ant brain. *Current Biology*, 26, 391–395. <https://doi.org/10.1016/j.cub.2015.12.040>.
- Linn, M., Glaser, S. M., Peng, T., & Grüter, C. (2020). Octopamine and dopamine mediate waggle dance following and information use in honeybees. *Proceedings of the Royal Society B*, 287, 20201950. <https://doi.org/10.1098/rspb.2020.1950>.
- Love, M., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15, 1165–1186. <https://doi.org/10.1186/s13059-014-0550-8>.
- Maboudi, M., Amini, J., Hahn, M., & Saati, M. (2017). Object-based road extraction from satellite images using ant colony optimization. *International Journal of Remote Sensing*, 38, 179–198. <https://doi.org/10.1080/01431161.2016.1264026>.
- McKenzie, S. K., Oxley, P. R., & Kronauer, D. J. (2014). Comparative genomics and transcriptomics in ants provide new insights into the evolution and function of odorant binding and chemosensory proteins. *BMC Genomics*, 15, 718. <https://doi.org/10.1186/1471-2164-15-718>.
- McQuillan, H. J., Barron, A. B., & Mercer, A. R. (2012). Age-and behaviour-related changes in the expression of biogenic amine receptor genes in the antennae of honey bees (*Apis mellifera*). *Journal of Comparative Physiology A*, 198, 753–761. <https://doi.org/10.1007/s00359-012-0745-y>.
- Menzel, R., & Greggers, U. (2013). Guidance by odors in honeybee navigation. *Journal of Comparative Physiology A Neuroethology Sensory Neural and Behavioral Physiology*, 199, 867–873. <https://doi.org/10.1007/s00359-013-0850-6>.
- Mercer, A. R., & Menzel, R. (1982). The effects of biogenic amines on conditioned and unconditioned responses to olfactory stimuli in the honeybee *Apis mellifera*. *Journal of Comparative Physiology*, 145, 363–368. <https://doi.org/10.1007/BF00619340>.
- Messan, M. R., Page, R. E. Jr, & Kang, Y. (2018). Effects of vitellogenin in age polyethism and population dynamics of honeybees. *Ecological Modelling*, 388, 88–107. <https://doi.org/10.1016/j.ecolmodel.2018.09.011>.
- Möglich, M., Maschwitz, U., & Hölldobler, B. (1974). Tandem calling: A new kind of signal in ant communication. *Science*, 186, 1046–1047. <https://doi.org/10.1126/science.186.4168.1046>.
- Morandin, C., Havukainen, H., Kulmuni, J., Dhaygude, K., Trontti, K., & Helanterä, H. (2014). Not only for egg yolk—functional and evolutionary insights from expression, selection, and structural analyses of *Formica* ant vitellogenins. *Molecular Biology and Evolution*, 31, 2181–2193. <https://doi.org/10.1093/molbev/msu171>.
- Mustard, J. A., Kurshan, P. T., Hamilton, I. S., Blenau, W., & Mercer, A. R. (2005). Developmental expression of a tyramine receptor gene in the brain of the honey bee, *Apis mellifera*. *Journal of Comparative Neurology*, 483, 66–75. <https://doi.org/10.1002/cne.20420>.
- Nagari, M., & Bloch, G. (2012). The involvement of the antennae in mediating the brood influence on circadian rhythms in “nurse” honey bee (*Apis mellifera*) workers. *Journal of Insect Physiology*, 58, 1096–1103. <https://doi.org/10.1016/j.jinsphys.2012.05.007>.
- Nelson, C. M., Ihle, K. E., Fondré, M. K., Page, R. E., & Amdam, G. V. (2007). The gene *vitellogenin* has multiple coordinating effects on social organization. *PLoS Biology*, 5, e86. <https://doi.org/10.1371/journal.pbio.0050062>.
- Nunes, F. M. F., Ihle, K. E., Mutti, N. S., Simoes, Z. L. P., & Amdam, G. V. (2013). The gene *vitellogenin* affects microRNA regulation in honey bee (*Apis mellifera*) fat body and brain. *Journal of Experimental Biology*, 216, 3724–3732. <https://doi.org/10.1242/jeb.089243>.
- Ogata, H., Goto, S., Sato, K., Fujibuchi, W., Bono, H., & Kanehisa, M. (1999). KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Research*, 27(1), 29–34. <https://doi.org/10.1093/nar/27.1.29>.
- Ozaki, M., Wada-Katsumata, A., Fujikawa, K., Iwasaki, M., Yokohari, F., Satoh, Y., & Yamaoka, R. (2005). Ant nestmate and non-nestmate discrimination by a chemosensory sensillum. *Science*, 309, 311–314. <https://doi.org/10.1126/science.1105244>.
- Pankiw, T., Roman, R., Sagili, R. R., & Zhu-Salzman, K. (2004). Pheromone-modulated behavioral suites influence colony growth in the honey bee (*Apis mellifera*). *Naturwissenschaften*, 91(12), 575–578. <https://doi.org/10.1007/s00114-004-0568-y>.
- Pantano, L. (2019). DEReport: Report of DEG analysis.
- Paoli, M., Weisz, N., Antolini, R., & Haase, A. (2016). Spatially resolved time-frequency analysis of odour coding in the insect antennal lobe. *European Journal of Neuroscience*, 44, 2387–2395. <https://doi.org/10.1111/ejn.13344>.
- Pelosi, P., Calvello, M., & Ban, L. (2005). Diversity of odorant-binding proteins and chemosensory proteins in insects. *Chemical Senses*, 30, i291–i292. <https://doi.org/10.1093/chemse/bjh229>.
- Peng, T., Derstroff, D., Maus, L., Bauer, T., & Grüter, C. (2021). Forager age and foraging state, but not cumulative foraging activity, affect biogenic amine receptor gene expression in the honeybee mushroom bodies. *Genes, Brain and Behavior*, 20, e12722. <https://doi.org/10.1111/gbb.12722>.
- Peng, T., Schroeder, M., & Grüter, C. (2020). Octopamine increases individual and collective foraging in a neotropical stingless bee. *Biology Letters*, 16, 20200238. <https://doi.org/10.1098/rsbl.2020.0238>.
- Perry, C. J., & Barron, A. B. (2013). Neural mechanisms of reward in insects. *Annual Review of Entomology*, 58, 543–562. <https://doi.org/10.1146/annurev-ento-120811-153631>.
- R Core Team (2018). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing.
- Reim, T., & Scheiner, R. (2014). Division of labour in honey bees: Age- and task-related changes in the expression of octopamine receptor genes. *Insect Molecular Biology*, 23, 833–841. <https://doi.org/10.1111imb.12130>.
- Reinhard, J., & Srinivasan, M. V. (2009). The role of scents in honey bee foraging and recruitment. In S. Jarau, & M. Hrncir (Eds.), *Food exploitation by social insects: Ecological, behavioral, and theoretical approaches* (pp. 165–182). CRC Press.
- Reinhard, J., Srinivasan, M. V., & Zhang, S. (2004). Scent-triggered navigation in honeybees. *Nature*, 427, 411. <https://doi.org/10.1038/427411a>.
- Rendell, L., Boyd, R., Cownden, D., Enquist, M., Eriksson, K., Feldman, M. W., Fogarty, L., Ghirlanda, S., Lillicrap, T., & Laland, K. N. (2010). Why copy others? Insights from the social learning strategies tournament. *Science*, 328, 208–213. <https://doi.org/10.1126/science.1184719>.
- Richter, M. R., & Waddington, K. D. (1993). Past foraging experience influences honey bee dance behaviour. *Animal Behaviour*, 46, 123–128. <https://doi.org/10.1006/anbe.1993.1167>.

- Rieucau, G., & Giraldeau, L. A. (2011). Exploring the costs and benefits of social information use: An appraisal of current experimental evidence. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 366, 949–957. <https://doi.org/10.1098/rstb.2010.0325>.
- Robertson, H. M., & Wanner, K. W. (2006). The chemoreceptor superfamily in the honey bee, *Apis mellifera*: Expansion of the odorant, but not gustatory, receptor family. *Genome Research*, 16, 1395–1403. <https://doi.org/10.1101/gr.5057506>.
- Robinson, G. E., Fernald, R. D., & Clayton, D. F. (2008). Genes and social behavior. *Science*, 322, 896–900. <https://doi.org/10.1126/science.1159277>.
- Rogers, L. J., & Vallortigara, G. (2008). From antenna to antenna: Lateral shift of olfactory memory recall by honeybees. *PLoS One*, 3, e2340. <https://doi.org/10.1371/journal.pone.0002340>.
- Sarma, M. S., Rodriguez-Zas, S. L., Hong, F., Zhong, S., & Robinson, G. E. (2009). Transcriptomic profiling of central nervous system regions in three species of honey bee during dance communication behavior. *PLoS One*, 4, e6408. <https://doi.org/10.1371/journal.pone.0006408>.
- Sasaki, K., Matsuyama, S., Harano, K. I., & Nagao, T. (2012). Caste differences in dopamine-related substances and dopamine supply in the brains of honeybees (*Apis mellifera* L.). *General and Comparative Endocrinology*, 178, 46–53. <https://doi.org/10.1016/j.ygcen.2012.04.006>.
- Scheiner, R., & Arnold, G. (2010). Effects of patriline on gustatory responsiveness and olfactory learning in honey bees. *Apidologie*, 41, 29–37. <https://doi.org/10.1051/apido/2009040>.
- Scheiner, R., Entler, B. V., Barron, A. B., Scholl, C., & Thamm, M. (2017). The effects of fat body tyramine level on gustatory responsiveness of honeybees (*Apis mellifera*) differ between behavioral castes. *Frontiers in Systems Neuroscience*, 11, 55. <https://doi.org/10.3389/fnins.2017.00055>.
- Scheiner, R., Erber, J., & Page, R. E. Jr (1999). Tactile learning and the individual evaluation of the reward in honey bees (*Apis mellifera* L.). *Journal of Comparative Physiology A*, 185, 1–10. <https://doi.org/10.1007/s003590050360>.
- Scheiner, R., Kuritz-Kaiser, A., Menzel, R., & Erber, J. (2005). Sensory responsiveness and the effects of equal subjective rewards on tactile learning and memory of honeybees. *Learning & Memory*, 12, 626–635. <https://doi.org/10.1101/lm.98105>.
- Scheiner, R., Page, R. E. Jr, & Erber, J. (2001). The effects of genotype, foraging role, and sucrose responsiveness on the tactile learning performance of honey bees (*Apis mellifera* L.). *Neurobiology of Learning and Memory*, 76, 138–150. <https://doi.org/10.1006/nlme.2000.3996>.
- Scheiner, R., Plückhahn, S., Öney, B., Blenau, W., & Erber, J. (2002). Behavioural pharmacology of octopamine, tyramine and dopamine in honey bees. *Behavioural Brain Research*, 136, 545–553. [https://doi.org/10.1016/S0166-4328\(02\)00205-X](https://doi.org/10.1016/S0166-4328(02)00205-X).
- Scheiner, R., Reim, T., Søvik, E., Entler, B. V., Barron, A. B., & Thamm, M. (2017). Learning, gustatory responsiveness and tyramine differences across nurse and forager honeybees. *Journal of Experimental Biology*, 220(8), 1443–1450. <https://doi.org/10.1242/jeb.152496>.
- Schulz, D. J., Elekonich, M. M., & Robinson, G. E. (2003). Biogenic amines in the antennal lobes and the initiation and maintenance of foraging behavior in honey bees. *Journal of Neurobiology*, 54, 406–416. <https://doi.org/10.1002/neu.10138>.
- Schulz, D. J., & Robinson, G. E. (1999). Biogenic amines and division of labor in honey bee colonies: Behaviorally related changes in the antennal lobes and age-related changes in the mushroom bodies. *Journal of Comparative Physiology A*, 184, 481–488. <https://doi.org/10.1007/s003590050348>.
- Sharma, K. R., Enzmann, B. L., Schmidt, Y., Moore, D., Jones, G. R., Parker, J., Berger, S. L., Reinberg, D., Zwiebel, L. J., Breit, B., Liebig, J., & Ray, A. (2015). Cuticular hydrocarbon pheromones for social behavior and their coding in the ant antenna. *Cell Reports*, 12, 1261–1271. <https://doi.org/10.1016/j.celrep.2015.07.031>.
- Sinakevitch, I., Niwa, M., & Strausfeld, N. J. (2005). Octopamine-like immunoreactivity in the honey bee and cockroach: Comparable organization in the brain and subesophageal ganglion. *Journal of Comparative Neurology*, 488, 233–254. <https://doi.org/10.1002/cne.20572>.
- Strassmann, J. (2001). The rarity of multiple mating by females in the social Hymenoptera. *Insectes Sociaux*, 48, 1–13. <https://doi.org/10.1007/PL00001737>.
- Strausfeld, N. J., Sinakevitch, I., Brown, S. M., & Farris, S. M. (2009). Ground plan of the insect mushroom body: Functional and evolutionary implications. *Journal of Comparative Neurology*, 513, 265–291. <https://doi.org/10.1002/cne.21948>.
- Taborsky, B., & Oliveira, R. F. (2012). Social competence: An evolutionary approach. *Trends in Ecology & Evolution*, 27, 679–688. <https://doi.org/10.1016/j.tree.2012.09.003>.
- Thamm, M., Scholl, C., Reim, T., Grübel, K., Möller, K., Rössler, W., & Scheiner, R. (2017). Neuronal distribution of tyramine and the tyramine receptor AmTAR1 in the honeybee brain. *Journal of Comparative Neurology*, 525, 2615–2631. <https://doi.org/10.1002/cne.24228>.
- Thom, C., Gilley, D. C., Hooper, J., & Esch, H. E. (2007). The scent of the waggle dance. *PLoS Biology*, 5, e228. <https://doi.org/10.1371/journal.pbio.0050228>.
- Tofilski, A. (2009). Shorter-lived workers start foraging earlier. *Insectes Sociaux*, 56, 359–366. <https://doi.org/10.1007/s00040-009-0031-3>.
- Toth, A. L., & Robinson, G. E. (2009). Evo-devo and the evolution of social behavior: Brain gene expression analyses in social insects. *Cold Spring Harbor Symposia on Quantitative Biology*, 74, 419–426. <https://doi.org/10.1101/sqb.2009.74.026>.
- Toth, A. L., Varala, K., Henshaw, M. T., Rodriguez-Zas, S. L., Hudson, M. E., & Robinson, G. E. (2010). Brain transcriptomic analysis in paper wasps identifies genes associated with behaviour across social insect lineages. *Proceedings of the Royal Society B: Biological Sciences*, 277, 2139–2148. <https://doi.org/10.1098/rspb.2010.0090>.
- van Zweden, J. S., & d'Ettorre, P. (2010). Nestmate recognition in social insects and the role of hydrocarbons. In G. J. Blomquist (Ed.), *Insect hydrocarbons: Biology, biochemistry and chemical ecology* (pp. 222–243). Cambridge University Press.
- Vavricka, C. J., Han, Q., Mehere, P., Ding, H., Christensen, B. M., & Li, J. (2014). Tyrosine metabolic enzymes from insects and mammals: A comparative perspective. *Insect Science*, 21, 13–19. <https://doi.org/10.1111/1744-7917.12038>.
- Vergoz, V., McQuillan, H. J., Geddes, L. H., Pullar, K., Nicholson, B. J., Paulin, M. G., & Mercer, A. R. (2009). Peripheral modulation of worker bee responses to queen mandibular pheromone. *Proceedings of the National Academy of Sciences*, 106, 20930–20935. <https://doi.org/10.1073/pnas.0907563106>.
- Vergoz, V., Roussel, E., Sandoz, J. C., & Giurfa, M. (2007). Aversive learning in honeybees revealed by the olfactory conditioning of the sting extension reflex. *PLoS One*, 2, e288. <https://doi.org/10.1371/journal.pone.0000288>.
- Voigt, J. P., & Fink, H. (2015). Serotonin controlling feeding and satiety. *Behavioural Brain Research*, 277, 14–31. <https://doi.org/10.1016/j.bbr.2014.08.065>.
- Von Frisch, K. (1967). *The dance language and orientation of bees*. Harvard University Press.
- Wallberg, A., Bunikis, I., Pettersson, O. V., Mosbech, M.-B., Childers, A. K., Evans, J. D., Mikheyev, A. S., Robertson, H. M., Robinson, G. E., & Webster, M. T. (2019). A hybrid de novo genome assembly of the honeybee, *Apis mellifera*, with chromosome-length scaffolds. *BMC Genomics*, 20, 275. <https://doi.org/10.1186/s12864-019-5642-0>.
- Wegener, J., Lorenz, M. W., & Bienenfeld, K. (2009). Physiological consequences of prolonged nursing in the honey bee. *Insectes Sociaux*, 56, 85–93. <https://doi.org/10.1007/s00040-008-1042-1>.

- Weimerskirch, H., Bertrand, S., Silva, J., Marques, J. C., & Goya, E. (2010). Use of social information in seabirds: Compass rafts indicate the heading of food patches. *PLoS One*, 5, e9928. <https://doi.org/10.1371/journal.pone.0009928>.
- Whitfield, C. W., Cziko, A. M., & Robinson, G. E. (2003). Gene expression profiles in the brain predict behavior in individual honey bees. *Science*, 302, 296–299. <https://doi.org/10.1126/science.1086807>.
- Woyciechowski, M., & Moroń, D. (2009). Life expectancy and onset of foraging in the honeybee (*Apis mellifera*). *Insectes Sociaux*, 56, 193–201. <https://doi.org/10.1007/s00040-009-0012-6>.
- Wray, M. K., Klein, B. A., & Seeley, T. D. (2012). Honey bees use social information in waggle dances more fully when foraging errors are more costly. *Behavioral Ecology*, 23, 125–131. <https://doi.org/10.1093/beheco/arr165>.
- Zayed, A., & Robinson, G. E. (2012). Understanding the relationship between brain gene expression and social behavior: Lessons from the honey bee. *Annual Review of Genetics*, 46, 591–615. <https://doi.org/10.1146/annurev-genet-110711-155517>.
- Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects models and extensions in ecology with R*. Springer.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Kennedy A, Peng T, Glaser SM, Linn M, Foitzik S, Grüter C. Use of waggle dance information in honey bees is linked to gene expression in the antennae, but not in the brain. *Mol Ecol*. 2021;00:1–13. <https://doi.org/10.1111/mec.15893>